

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE		2. REPORT TYPE Viewgraphs		3. DATES COVERED	
4. TITLE AND SUBTITLE Polarization Properties of Chiral-Core Planar Waveguides		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S) Dr. Warren N. Herman		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Air Warfare Center Aircraft Division 22347 Cedar Point Road, Unit #6 Patuxent River, Maryland 20670-1161		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT A new form for the modal eigenvalue equations of chiral-core planar waveguides provides insight into the transition, with increasing chirality, from TE/TM (transverse electric/magnetic) modes in the archiral case to nearly right-handed and left-handed circularly polarized modes. Dramatic variation of the polarization eccentricity with thickness and frequency is discussed.					
15. SUBJECT TERMS Polymers; Chiral; waveguides					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON Dr. Warren Herman
a. REPORT	b. ABSTRACT	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code) (301) 342-9114

Standard Form 298 (Rev. 8-98)
Prescribed by ANSI Std. Z39-18

20010824 082

Polarization properties of chiral-core planar waveguides

Presenter: Warren N. Herman

Address: US Navy, NAWCAD
EO Sensors Division AIR456
Patuxent River, MD 20670

Abstract

A new form for the modal eigenvalue equations of chiral-core planar waveguides provides insight into the transition, with increasing chirality, from TE/TM (transverse electric/magnetic) modes in the achiral case to nearly right-handed and left-handed circularly polarized modes. Dramatic variation of the polarization eccentricity with thickness and frequency is discussed.




Polarization Properties of Chiral Core Planar Waveguides

W.N. Herman
U.S. Navy, NAWCAD
EO Sensors Division, Patuxent River, MD

OTF2001
15-17 October 2001



Chiral Core Optical Waveguides



ISOTROPIC CHIRAL MEDIUM

Constitutive Relations
(Drude-Born-Federov)

$$\vec{D} = \epsilon(\vec{E} + \gamma \nabla \times \vec{E}) \quad \vec{B} = \mu(\vec{H} + \gamma \nabla \times \vec{H})$$

Bohren's Decomposition

$$\vec{E} = \frac{1}{2}(\vec{F}^+ + \vec{F}^-) \quad \vec{H} = \frac{1}{2i} \sqrt{\frac{\epsilon}{\mu}}(\vec{F}^+ - \vec{F}^-)$$

Wave Equation

$$\nabla^2 \vec{F}^\pm + (k_0 n_\pm)^2 \vec{F}^\pm = 0$$

Eigenmodes in bulk
material circularly polarized

$$\vec{F}^\pm = \vec{E}_0 e^{i(k_0 n_\pm z - \omega t)} (\hat{x} \pm i\hat{y})$$

Refractive indices
for RH and LH
waves

$$n_\pm = \frac{n_g}{1 \pm \delta} \quad \delta = k_0 n_g \gamma \quad n_g = \sqrt{\frac{\epsilon}{\epsilon_0}} \quad k_0 = \frac{2\pi}{\lambda}$$

Rotatory Power


$$\rho = \frac{\pi(n_- - n_+)}{\lambda} \approx k_0 n_g \delta$$

A. Lahtakia, V.K. Varadan, and V.V. Varadan, Time-Harmonic Electromagnetic Fields in Chiral Media, Lecture Notes in Physics Series 335 (Springer-Verlag, Berlin, 1989).
I.V. Lindell, A.H. Sihvola, S.A. Tretyakov, A.J. Viitanen, Electromagnetic Waves in Chiral and Bi-isotropic Media, (Artech House, Norwood, MA, 1994).

Modes in Chiral Asymmetric Waveguide

W. N. Herman, accepted *J. Opt. Soc. Am. A*

$$\vec{F}^\pm(y, z) = \vec{\Psi}^\pm(y) \exp(-ik_0 n_{eff} z)$$



Modal equations: (3 equations to be solved simultaneously for n_{eff} , g , h)

$$u^\pm d = \cot^{-1} \left(\sigma_0^\pm \frac{r_0 \pm g}{1 \pm g} \right) + \cot^{-1} \left(\sigma_s^\pm \frac{r_s \pm h}{1 \pm h} \right) + m^\pm \pi$$

$$h(g, n_{eff}) = \frac{(Sr_0^+ - Sr_0^-) + (S_0^+ + S_0^-)g}{(Sr_0^+ + Sr_0^-) + (S_0^+ - S_0^-)g}$$

$$S_0^\pm \equiv \sigma_0^\pm \sin(u^\pm d) \mp \cos(u^\pm d)$$

$$Sr_0^\pm \equiv r_0 \sigma_0^\pm \sin(u^\pm d) \mp \cos(u^\pm d)$$

$$\sigma_0^\pm \equiv (1 \pm \delta) \frac{u^\pm}{v}, \quad \sigma_s^\pm \equiv (1 \pm \delta) \frac{u^\pm}{w}, \quad r_0 \equiv \frac{n_0^2}{n_g^2}, \quad r_s \equiv \frac{n_s^2}{n_g^2}$$



$$u^\pm \equiv k_0 \sqrt{n_t^2 - n_{eff}^2}, \quad v \equiv k_0 \sqrt{n_0^2 - n_{eff}^2}, \quad w \equiv k_0 \sqrt{n_s^2 - n_{eff}^2}$$



$$n^\pm = \frac{n_g}{1 \pm \delta}$$



Parameters g , h determine eccentricity of polarization ellipse for transverse E-field:


$$\left. \frac{E_x}{E_y} \right|_{y \geq 0} = i \frac{n_{eff}}{n_g} \frac{1}{g}$$

$$\left. \frac{E_y}{E_x} \right|_{y \leq -d} = i \frac{n_{eff}}{n_g} \frac{1}{h}$$









1. M.M. Green, et al., *Angew. Chem. Int. Ed.* **38**, 3138(1999)
2. J.F. Nye, *Physical Properties of Crystals*(Oxford, 1960), p. 262.
3. T.J. Katz, et al., *J. Am. Chem. Soc.* **115**, 3182(1993)
4. C. Nuckolls, et al., *J. Am. Chem. Soc.* **121**, 79(1999)